



Experimental warming differentially affects microbial structure and activity in two contrasted moisture sites in a *Sphagnum*-dominated peatland



Frédéric Delarue^{a,b,c,*}, Alexandre Buttler^{d,e,f}, Luca Bragazza^{e,f,g}, Laurent Grasset^h, Vincent E.J. Jassey^{e,f}, Sébastien Gogo^{a,b,c}, Fatima Laggoun-Défarge^{a,b,c}

^a Université d'Orléans, ISTO, UMR 7327, 45071 Orléans, France

^b CNRS, ISTO, UMR 7327, 45071 Orléans, France

^c BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France

^d Laboratoire de Chrono-Environnement, UMR CNRS 6249, UFR des Sciences et Techniques, 16 route de Gray, Université de Franche-Comté, F-25030 Besançon, France

^e École Polytechnique Fédérale de Lausanne (EPFL), School of Architecture, Civil and Environmental Engineering (ENAC), Laboratory of Ecological Systems (ECOS), Bâtiment GR, Station 2, CH-1015 Lausanne, Switzerland

^f WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Site Lausanne, Station 2, Case postale 96, CH-1015 Lausanne, Switzerland

^g University of Ferrara, Department of Life Science and Biotechnology, Corso Ercole I d'Este 32, I-44121 Ferrara, Italy

^h CNRS, Laboratoire de Synthèse et de Réactivité des Substances Naturelles—UMR 6514, Université de Poitiers, 4 rue M. Brunet, 86022 Poitiers Cedex, France

HIGHLIGHTS

- Open-Top Chambers were used to raise air temperature by up to 1 °C in a peatland.
- Interaction between peat temperature and moisture content was investigated.
- Impact on peat decomposition was assessed by combining various microbial proxies.
- The effect of air warming occurred when comparing distinct moisture sites.
- We describe a change in microbial structure and enzymatic activities.

ARTICLE INFO

Article history:

Received 24 September 2014

Received in revised form 23 December 2014

Accepted 28 December 2014

Available online xxxxx

Editor: Mark Hanson

Keywords:

Carbon

Climate change

Enzymes

Microbial biomass

Moisture

Open-top chambers

ABSTRACT

Several studies on the impact of climate warming have indicated that peat decomposition/mineralization will be enhanced. Most of these studies deal with the impact of experimental warming during summer when prevalent abiotic conditions are favorable to decomposition. Here, we investigated the effect of experimental air warming by open-top chambers (OTCs) on water-extractable organic matter (WEOM), microbial biomasses and enzymatic activities in two contrasted moisture sites named Bog and Fen sites, the latter considered as the wetter ones. While no or few changes in peat temperature and water content appeared under the overall effect of OTCs, we observed that air warming smoothed water content differences and led to a decrease in mean peat temperature at the warmed Bog sites. This thermal discrepancy between the two sites led to contrasting changes in microbial structure and activities: a rise in hydrolytic activity at the warmed Bog sites and a relative enhancement of bacterial biomass at the warmed Fen sites. These features were not associated with any change in WEOM properties namely carbon and sugar contents and aromaticity, suggesting that air warming did not trigger any shift in OM decomposition. Using various tools, we show that the use of single indicators of OM decomposition can lead to fallacious conclusions.

Lastly, these patterns may change seasonally as a consequence of complex interactions between groundwater level and air warming, suggesting the need to improve our knowledge using a high time-resolution approach.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

The impact of climate change and its consequences on global air temperature is still debated as some uncertainties remain (IPCC, 2007; Otto et al., 2013), in particular concerning the potential feedback of

* Corresponding author at: Institut de Minéralogie, de Physique des Matériaux, et de Cosmochimie (IMPMC) Sorbonne Universités – Muséum National d'Histoire Naturelle, UPMC Univ Paris 06, UMR CNRS 7590, IRD UMR 206, 61 rue Buffon, F-75005 Paris, France.
E-mail address: fdelarue@mnhn.fr (F. Delarue).

the terrestrial carbon (C) cycle to climate warming (Davidson and Janssens, 2006; Friedlingstein et al., 2006). Understanding the fate of the C stored in peatlands is crucial since these ecosystems contain about one-third of the world's soil organic C as peat (Gorham, 1991), the equivalent of about 60% of atmospheric C.

The C sink function of peatlands is mainly the result of persistent anoxic conditions, low temperature and water acidity that reduce microbial decomposition and promote the accumulation of organic matter (OM) as peat. As a result, peat organic C is expected to be particularly sensitive to climate warming because of the higher intrinsic temperature sensitivity of this type of organic soil (Davidson and Janssens, 2006). Thus, a 1 °C increase in air temperature has been estimated to enhance C fluxes from heterotrophic respiration in northern peatlands by about 38–100 Mt of C per year (Dorrepaal et al., 2009). Other studies indicated that climate change can further diminish C sequestration by promoting the growth of vascular plants which, in turn, depress the productivity of peat mosses (Breeuwer et al., 2009; Bragazza et al., 2013). In contrast to these studies, Loisel and Yu (2013) and Charman et al. (2013) indicated that the C accumulation rate of many northern peatlands could increase in response to a warmer climate in the future, as long as moisture is not a limiting factor. These examples illustrate the ongoing debate on the fate of C in peatlands in response to climate change. Several studies demonstrated the complex interaction between peat moisture and air temperature in regulating peat decomposition (Delarue et al., 2011a; Jassey et al., 2011; Bokhorst et al., 2013) but few experimental studies have specifically addressed this topic, notwithstanding the close relationship between moisture, temperature and C cycling for peat decomposition (McNeil and Waddington, 2003). It was demonstrated that an increased water evaporation in peat soils and a drying out of the surface layer can decrease the soil's thermal conductivity which, in turn, can prevent heat from propagating deeper into the soil and can therefore keep them colder (Dabros and Fyles, 2010). Other authors also reported that evaporation was associated with a cooling of the upper moss layers, inducing a condensation of vapour (Carleton and Dunham, 2003).

The peculiar environmental conditions in peatlands favour the establishment of *Sphagnum* mosses, which are known to produce recalcitrant litters enriched in polyphenolic compounds (van Breemen, 1995; Abbott et al., 2013; Swain and Abbott, 2013). Although polyphenols inhibit extracellular enzymatic activity (Freeman et al., 2001, 2004), the enzymes belonging to the phenoloxidase (PO) group have the ability to degrade recalcitrant polyphenols accumulating in peatlands (McLatchey and Reddy, 1998; Freeman et al., 2001). In a perspective of climate change, PO activity is expected to increase as a consequence of more frequent drought events and associated oxygenation of peat soils (Fenner and Freeman, 2011). Due to a decrease in soluble phenols with increasing activity of PO, hydrolytic enzymes such as leucine amino-peptidase (LAP), β -glucosidase (BG) and acidic phosphatase (AP) are no longer inhibited so that the breakdown of OM can start. Such a pattern is known as the enzymatic latch theory (Freeman et al., 2001). Following such cascading effects, various studies suggested that peat OM decomposition will be enhanced by climate warming (Dorrepaal et al., 2009; Fenner and Freeman, 2011; Jassey et al., 2013). However, the studies were mainly conducted during the summer months, when environmental constraints, i.e. water level drawdown and air temperature, were less limiting for microbial metabolism. Therefore, two major questions remain to be clarified: (1) how air temperature and water level interact to affect both soil temperature and moisture under conditions of water saturated peat and (2) how this interaction can affect the soil C cycle.

In this study we investigate the interactive effects of air warming which was experimentally induced by Open-Top Chambers (OTCs), and water level by comparing two habitats characterized by contrasted soil moisture conditions. The study was performed in early summer when both soil water level and air temperature had not yet reached their annual minimum and maximum values respectively. Specifically,

we explored peat moisture changes as a function of soil water level, and air and peat temperatures. The impact of air warming and peat moisture on the soil C cycle was assessed using phospholipid fatty acids as an index of microbial biomass, various enzymatic activities (leucine aminopeptidase, β -glucosidase, acidic phosphatase and phenol oxidase) as indexes of microorganism activities and the corresponding chemical quality of water-extractable OM (WEOM).

2. Material and methods

2.1. Study site, experimental design and sampling

The study site is an undisturbed ombrotrophic *Sphagnum*-dominated peatland situated in the Jura Mountains (Le Forbonnet, France; 46°49'35"N, 6°10'20"E), at an altitude of ca. 840 m a.s.l. The annual mean temperature at the site is ca. 6.5 °C, and the annual precipitation is about 1200 mm (Delarue et al., 2011b). Cold winters (mean monthly temperature ca. 1.4 °C) and mild summers (ca. 14.6 °C) characterize the climate.

Peat samples were collected in late June 2011 across a vegetation gradient corresponding to a narrow transitional minerotrophic to ombrotrophic area. The transition from the minerotrophic to the ombrotrophic area was characterized by a shift from an area of relatively flat and homogeneous surface dominated by *Sphagnum fallax* with a low abundance of vascular plants (i.e. *Eriophorum vaginatum*, *Vaccinium oxycoccus* and *Andromeda polifolia*) to a surface with a patterned vegetation of hummocks, where *Sphagnum magellanicum*, *V. oxycoccus*, *E. vaginatum* and *Calluna vulgaris* developed, and hollows mainly occupied by *S. fallax*, *Carex rostrata* and *A. polifolia*. The main change between the minerotrophic and the ombrotrophic areas was the occurrence of *S. magellanicum* in the latter, entailing a change in the microtopography (considered as a site effect). For simplicity's sake, we will call the minerotrophic and ombrotrophic areas, respectively, Fen and Bog sites hereafter.

The experimental design was described in detail in previous work (Delarue et al., 2011a; Jassey et al., 2011). Briefly, OTCs are passive warming chambers designed following the International Tundra Experiment (ITEX) to obtain quasi-natural transmittance of visible wavelengths and to minimize the transmittance of re-radiated infrared wavelengths (Marion et al., 1997; Aronson and McNulty, 2009). The hexagonal chambers are made of transparent polycarbonate and are 50 cm high, 1.7 m wide at the top and 2.4 m wide at the base. They were raised 10 cm above the soil surface to allow air to circulate. Six plots were equipped with OTCs in May 2008, and 6 other plots were used as controls (CTLs). For this study, the plots were named as follows: Bog-OTC and Bog-CTL for plots in the Bog site with and without OTCs respectively, and Fen-OTC and Fen-CTL for plots in the Fen site with and without OTCs respectively.

Temperature of the peat (7 cm deep) and air (10 cm above *Sphagnum* capitulum) was automatically measured every 30 min using thermocouple probes in each plot and a data logger (CR-1000 Campbell). Monthly mean, minimum and maximum temperatures, for both peat and air, were then calculated for the period from January 2011 to June 2011. The ground water level was automatically measured in one randomly selected plot at both Bog and Fen sites (mid-May 2011 to late June 2011). Lastly, peat moisture and temperature were measured at ca. 5 cm depth by Decagon® sensors only during the growing season (from early May 2011 to October 2011) in two randomly selected plots at both the Bog and Fen sites. Twelve peat cores 30 cm long were sampled in June 2011, after 3 years of experimental warming. The peat cores were cut into five slices (0 to 5, 5 to 10, 10 to 15, 15 to 20 and 20–25 cm interval depth) and frozen. Within two weeks after sampling, each slice was subdivided into two parts. For each part, water was gently extracted following the procedure described by Delarue et al. (2011a) modified by the use of a PTFE filter (0.45 μ m pore size). After WEOM extraction, peat samples were dried at 105 °C for 24 h in order to obtain the peat dry mass. The

water content was calculated by considering the peat dry mass and the peat wet mass measured before WEOM extraction.

2.2. Structure of microbial communities – PLFAs

Phospholipid-fatty acids (PLFAs) were extracted on freeze-dried peat samples using the method of Bligh and Dyer (1959), modified for peat (Andersen et al., 2010). Peat samples (250 mg) were shaken for 2 h in a phosphate-buffer (0.1 M; pH 7):CHCl₃:MeOH (0.9:1:2 v/v/v) solution and the supernatant was then transferred to a phosphate buffer/chloroform solution. After separation of the organic phase overnight, the CHCl₃-lipid phase was split into neutral, glyco- and phospholipids in a silicic acid column by eluting chloroform, acetone and methanol respectively. Phospholipids were then transesterified into fatty acid methyl esters (FAMES) after incubation at 40 °C in a methanolic KOH (1 M):toluene (1:1 v/v) solution. The solution was neutralized with acetic acid (1 M) and FAMES were extracted by adding a hexane:CHCl₃ (4:1) solution. The hexane fraction was then passed through an MgSO₄ column before evaporation to dryness under an N₂ flux.

FAMES were analysed by means of GC-MS and quantified using a GC apparatus (Trace GC, Thermo Finnigan) equipped with a Supelco Equity 5-fused silica column (30 m length, 0.25 mm internal diameter, 0.25 µm film thickness) coupled to a mass spectrometer (Quadrupole DSI II, Thermo Finnigan). Helium was employed as the carrier gas at a constant flow rate. Methyl nonadecanoate (C₁₉O₂Me) was used as internal standard. Strict location of double bonds was realized by derivatization of FAMES into picolinyl esters on representative samples (Wretensjö et al., 1990).

We used the PLFAs i15:0, a15:0, i16:0, i17:0 and a17:0 as markers of G+ bacteria (Frostegård and Bååth, 1996); 16:1ω7c and cy17:0 as markers of G- bacteria (Wilkinson, 1988; Zelles, 1999); 18:2ω6,9 as marker of Fungi (Bardgett et al., 1996; Frostegård and Bååth, 1996; Zelles, 1999); 10Me16:0 and 10Me18:0 as markers of actinobacteria and sulfate-reducing bacteria (Kroppenstedt, 1985) and 20:4ω6,9,12,15 as markers of Protozoa (Ringelberg et al., 1997). Other PLFAs detected in the samples were not specific to one particular functional group so were not used in the comparisons. PLFA concentrations are expressed as µg C·g⁻¹ of dry peat.

2.3. Water-extractable organic matter analyses

WEOM analyses were performed on the first part of each peat slice. The WEOM was divided into three aliquots for analyses of organic C (WEOC), total sugars and SUVA₂₈₀, an index of the aromaticity of WEOM (Kalbitz et al., 2003). To calculate the WEOC, the dissolved organic carbon (DOC in mg l⁻¹) was first determined after acidification with H₃PO₄ (pH = 4) and N₂ purging. DOC was then measured with a Shimadzu SSM-5000A total carbon analyser. Finally, the mass of dissolved C was calculated and divided by the initial sample dry mass to obtain the WEOC expressed in mg·g⁻¹ of dry peat. Total sugars were determined on the second aliquot following the phenol-sulfuric acid method with glucose as standard to allow the calculation of sugar content (Dubois et al., 1956). Total sugar contents were expressed in mg of carbon g⁻¹ of dry peat, since we assumed that the weight ratio of C in sugars was that of glucose (2.5). For SUVA₂₈₀, the third aliquot was adjusted to a pH ranging from 6 to 7 following the recommendation of Weishaar et al. (2003). UV absorbance was then measured at 280 nm using a UV spectrophotometer. Finally, SUVA₂₈₀ was calculated as absorbance divided by WEOC concentration (Hansson et al., 2010) and is expressed as g of dry peat per mg C⁻¹ cm⁻¹.

2.4. Extracellular enzymatic assays

Enzymatic activities were measured on the second subsample of each slice. The activity of extracellular phenol oxidase was

Table 1

Results of a Repeated Measures ANOVA's to test the singular and overall interaction effects of time (n = 6), site (n = 6) and experimental warming (n = 6) on air and peat temperatures. Mean, minimum and maximum monthly temperatures from January 2011 to June 2011 were used as repeated measures. Significant differences are in bold and indicated by a p-value below 0.05.

Effect	df	Mean		Minimum		Maximum	
		F	p	F	p	F	p
<i>Air temperatures</i>							
Site	1	2.44	0.16	11.65	<0.05	5.77	<0.05
Treatment	1	42.94	<0.05	15.71	<0.05	16.02	<0.05
Time	5	25639.89	<0.05	4220.60	<0.05	3492.06	<0.05
Site × Treat.	1	0.36	0.56	1.78	0.22	0.23	0.64
Site × Time	5	4.23	<0.05	2.74	<0.05	5.09	<0.05
Treat. × Time	5	21.16	<0.05	1.26	0.30	15.20	<0.05
Site × Treat. × Time	5	0.29	0.92	0.38	0.86	0.19	0.97
<i>Peat temperatures</i>							
Site	1	2.81	0.13	0.01	0.93	1.77	0.22
Treatment	1	0.01	0.91	6.69	<0.05	2.01	0.19
Time	5	1389.12	<0.05	969.23	<0.05	391.76	<0.05
Site × Treat.	1	3.28	0.11	6.90	<0.05	0.01	0.92
Site × Time	5	1.37	0.26	1.01	0.42	1.14	0.35
Treat. × Time	5	1.03	0.41	6.08	<0.05	2.01	0.10
Site × Treat. × Time	5	0.32	0.90	3.63	<0.05	0.28	0.92

determined spectrophotometrically by using 10 mM L-dopa (dihydroxyphenylalanine) solution as substrate (Pind et al., 1994). The activity of phenol oxidase (PO) was expressed in µmol of 2,3-dihydroindole-5,6-quinone-2-carboxylate (dicq) min⁻¹ g⁻¹ of dry peat.

The activity of extracellular hydrolytic enzymes was measured by adding 4-methylumbelliferyl-β-D-glucoside for β-glucosidase (BG), L-leucine-7-amido-4-methylcoumarinhydrochloride for leucine aminopeptidase (LAP) and 4-MUF-phosphate for the activity of acidic phosphatase (AP) to about 1 g of fresh soil. After incubation (1 h for BG, and LAP, and 45 min for AP), the fluorescence of the supernatant after centrifugation was measured on a microplate reader (BioTek Synergy MX) at 450-nm emission and 330-nm excitation wavelength. To quantify product release and account for quenching effects, a set of standards was prepared using methylumbelliferone (MUF) and 7-amino-4-methylcoumarin (MCU) mixed with peat extract (Freeman et al., 1995; Saiya-Cork et al., 2002). Hydrolytic enzyme activity was expressed as µmol of substrate (MUF) converted per minute and per gramme of dry peat.

2.5. Statistics

To study the impact of air warming upon water content, PLFAs, WEOM features and extracellular enzymatic activities resulting from each depth, slices were pooled in order to obtain an overall response for the 25 cm peat column in each plot. All statistical analyses were performed using XLSTAT software (Addinsoft®). Data were tested for normality using the Kolmogorov-Smirnov test and for homogeneity of variance using the Levene test. Data were log₁₀-transformed when non-normality and/or no homogeneity of the variance were found. Variations in air and peat temperatures were examined through Repeated Measures ANalysis Of VAriance (MANOVA) in order to test the singular impact and interactions of sites, air warming and time (i.e., months for air and peat temperatures). Following significant MANOVA tests (p-value <0.05), significant differences were determined with Fisher's LSD tests. Variations in water content, PLFAs, WEOM features and extracellular enzymatic activities were analysed using ANOVA (i) to test the overall impact of air warming on these variables, (ii) to test the impact of air warming within the Bog and Fen sites and (iii) to investigate the impact of air warming on the initial differences distinguishing the Bog and Fen sites.

Table 2

Air and peat temperatures (°C) according to sites (n = 6), warming treatment (n = 6) and their interactions (n = 3). Mean, minimum and maximum temperatures were calculated according to monthly temperatures from January 2011 to June 2011. Significant differences were determined with the Fisher's LSD test and are indicated in bold and by a p-value below 0.05.

	Air temperatures			Peat temperatures		
	Mean temperature	Min. temperature	Max. temperature	Mean temperature	Min. temperature	Max. temperature
<i>Site (n = 6)</i>						
Bog	6.2	−4.2	19.3	5.1	4.2	6.2
Fen	6.0	−3.8	17.9	5.5	4.1	7.2
p-Value	0.16	<0.05	<0.05	0.13	0.93	0.22
<i>Treatment (n = 6)</i>						
Control	5.7	−4.2	17.4	5.3	4.6	6.1
Warmed	6.5	−3.8	19.7	5.3	3.7	7.2
p-Value	<0.05	<0.05	<0.05	0.91	<0.05	0.19
<i>Site × Treatment (n = 3)</i>						
Bog-CTL	5.7	−4.5	18.3	5.3	5.0	5.7
Bog-OTC	6.6	−3.9	20.3	4.9	3.3	6.7
Fen-Control	5.6	−4.0	16.6	5.3	4.1	6.6
Fen-OTC	6.3	−3.7	19.2	5.7	4.2	7.8
<i>p-Value</i>						
Bog-CTL vs. Bog-OTC	<0.05	<0.05	<0.05	0.27	<0.05	0.38
Fen-CTL vs. Fen-OTC	<0.05	<0.05	<0.05	0.21	0.98	0.31
Bog-CTL vs. Fen-CTL	0.52	<0.05	0.08	0.93	0.09	0.41
Bog-OTC vs. Fen-OTC	0.16	0.18	0.21	<0.05	0.11	0.34

3. Results

3.1. Air and peat temperatures

Continuous measurements of air temperature during the period from January to June 2011 in both control and OTC plots indicated significant effects related to the site type, warming treatment and time (Table 1). Minimum air temperature was significantly higher at the Fen site (−3.8 °C) than at the Bog site (−4.2 °C; Table 2). Conversely, maximum air temperature was significantly higher at the Bog site (19.3 °C) than at the Fen site (17.9 °C). The single effect of air warming treatment also led to a rise in mean (+0.8 °C), minimum (+0.4 °C) and maximum (+2.3 °C) air temperatures (Table 2). More specifically, the experimental air warming treatment increased the mean (+0.9 °C), minimum (+0.6 °C) and maximum (+2 °C) air temperatures at the Bog site (Table 2). At the Fen site, OTCs were also associated with a rise in mean (+0.7 °C), minimum (+0.3 °C) and maximum (+2.6 °C) air temperatures (Table 2).

Few specific effects of site or of the experimental warming were recorded on peat temperatures at 7 cm depth (Table 1). With OTC treatment, the minimum peat temperature decreased by 0.9 °C (Table 2). Two significant differences were also observed due to the interaction between site and air warming: the minimum peat temperature was lower at the Bog-OTC (3.3 °C) than at the Bog-CTL plot (5.0 °C)

(Table 2), and the mean peat temperature was higher at the Fen-OTC (5.7 °C) than at the Bog-OTC plot (4.9 °C). A week before the sampling, the pattern was similar (Fig. 1), with no significant effect of experimental warming on peat temperature at the Fen site but a significant decrease in minimum peat temperature at the Bog site (from 11.1 to 10.4 °C). Additionally, experimental warming also induced a decrease of mean peat temperature in the Bog-OTC site (11.3 °C) as compared to the Fen-OTC (12.9 °C) site.

3.2. Ground water level and peat moisture changes

From mid-May to late June 2011, the ground water level was systematically higher at the Fen site than at the Bog site (+3 cm; Fig. 2) and it was strongly correlated with peat moisture from early May 2011 to late June 2011 (Fig. 3A). During this period, no significant relationship was found between peat moisture and air and peat temperatures (Fig. 3B and C). Instead, there was a positive correlation between air and peat temperatures ($p < 0.05$) at both the Bog and Fen sites (Fig. 3D).

No overall effect of warming treatment was observed on peat moisture (Table 3). More specifically, no significant changes were recorded at the Fen site, but water content was significantly higher in the Bog-OTC (94.5%) site than in the corresponding control plots (93.3%; Table 3). Peat moisture in the control plots at the Bog and Fen sites were not significantly different ($p = 0.07$), but this trend disappeared under the effect of air warming.




Fig. 1. Effect of experimental warming on mean, minimum and maximum peat temperatures at both Bog and Fen sites during the week before sampling. Each value corresponds to the weekly mean temperature, and to the minimum and maximum peat daily temperatures. Error bars are indicative of standard error between replicates (n = 3). Significant differences were tested using one-way ANOVA and are indicated by different letters.




Fig. 2. Ground water level (below the *Sphagnum capitulum*) measured at both Bog and Fen sites from mid-May to late June.




Fig. 3. Relationships between ground water level and peat moisture (A), peat moisture and air mean temperature (B), peat moisture and peat mean temperature (C), air and peat mean temperatures (D) at the Bog and Fen sites. Measurements were performed from mid-May 2011 to October 2011. Correlations were based on the Pearson's test. Each value corresponds to the daily mean ($n = 146$). Significant correlation is indicated by an asterisk ($p < 0.05$).

3.3. Water-extractable organic matter features and phospholipid fatty acids

No impact of air warming treatment was recorded on WEOC, sugar content and SUVA₂₈₀ (Table 3). With respect to PLFAs, significant changes occurred only when comparing the effect of air warming treatment in Bog and Fen sites (Table 3), while no significant differences were observed between their control sites. In warmed plots, PLFAs from G-positive and G-negative bacteria became significantly higher at the Fen site compared to the Bog site (respectively 47.3 and 18.8 $\mu\text{g C} \cdot \text{g}^{-1}$ of dry peat in the Fen site and 12.3 and 4.2 $\mu\text{g C} \cdot \text{g}^{-1}$ of dry peat in the Bog site – Table 3). The Fen control plots also had higher Protozoan contents as compared to the Bog control plots, but this difference did not persist under warming treatment.

3.4. Enzymatic activities

There was no overall effect of warming treatment on enzymatic activity (Table 3). Nevertheless, at the Bog site, warming treatment significantly enhanced AP activity. In warmed plots, the activity of LAP and AP was significantly higher at the Bog site as compared to the Fen site (respectively, 5.4 and 1.4 $\mu\text{mol MUF min}^{-1} \text{g}^{-1}$ at the Bog-site and 4.7 and 1.0 $\mu\text{mol MUF min}^{-1} \text{g}^{-1}$ at the Fen site).

4. Discussion

4.1. Experimental air warming enhances the discrepancy of peat temperatures between Bog and Fen sites

From January to June 2011, the OTCs enhanced mean air temperature up to 0.9 °C and 0.7 °C in the Bog and the Fen sites, respectively. Such a temperature rise is in accordance with other *in situ* warming experiments with OTCs (Sullivan et al., 2008; Dorrepaal et al., 2009; Weedon et al., 2012). The increase in air temperature was associated with a decrease in minimum peat temperature under the impact of warming, only at the Bog site (Tables 1 and 2). At the Bog site, the increase in air temperature was also associated with an increase in peat water content (Table 3). This result is surprising since most studies on

experimental warming reported a decrease or no effect on peat moisture (Hollister et al., 2006; Dorrepaal et al., 2009; Bokhorst et al., 2011; Delarue et al., 2011a; Jassey et al., 2013). The question is therefore whether this effect was due to an experimental artefact or whether it results from a thermodynamic constraint. It was demonstrated that OTCs can stop wind blowing, thus reducing evaporation (de Boeck et al., 2012). As no relationship was found between wind speed and peat moisture (at 5 cm depth) in the control plots (data not shown), we assume that no significant reduction of evaporation by OTCs occurred at 5 cm depth. However, we cannot rule out the effect of wind at the surface of the *Sphagnum* carpet. In both sites, peat moisture was mainly controlled by ground water level rather than by air temperature (Fig. 3A, B and C). Therefore, such a rise in peat moisture at the Bog site may result from an interaction between air temperature and ground water level owing to capillary strength. Water capillary flow is the main mass flux within peat (Price et al., 2009) so if the capillary flow is not strong enough to compensate for the evaporation rate, mosses start to dry out. Conversely, if the capillary flow compensates for the evaporation rate (Yazaki et al., 2006), then the vapour diffusion through evaporation can cool the upper peat layer (Carleton and Dunham, 2003). In addition, this can lead to the condensation of vapour in the upper peat layer, which causes a slight increase in peat moisture (Price et al., 2009). This mechanism can partially explain the observed increase in peat moisture and the decrease in minimum peat temperature at the Bog site. At the Fen site, no temperature changes were recorded. Due to the different effects of experimental warming in the two sites, it can be concluded that there is a thermal discrepancy, as indicated by the lower peat temperature at the Bog-OTC as compared to the Fen-OTC site (Table 2). Such a discrepancy was also measured the week before sampling (Fig. 1) and was associated with the disappearance of the peat moisture discrepancy between the warmed Bog and Fen sites (Table 3).

Overall, this discrepancy suggests that a slightly higher ground water level (about 3 cm; Fig. 2) may prevent any effect of experimental warming on both peat temperature and moisture, suggesting that a potential thermodynamic threshold occurs as a function of groundwater level.

Table 3

Effect of warming treatment on PLFAs (G+ bacteria, G– bacteria, Fungi, Actinobacteria and Protozoan in $\mu\text{g C g}^{-1}$ of dry peat), water content, WEOM characteristics (water extractable organic carbon in mg g^{-1} of dry peat, sugar content in mg g^{-1} of dry peat and SUVA₂₈₀ in $\text{g of dry peat per mg C}^{-1} \text{cm}^{-1}$) and extracellular enzymatic activities (phenoloxidase–PO in $\mu\text{mol of dicq min}^{-1} \text{g}^{-1}$ of dry peat, β -glucosidase–BG, leucine aminopeptidase–LA and acidic phosphatase–AP in $\mu\text{mol of substrate}$). The impact of warming was tested with Bog-CTL vs. Bog-OTC (n = 3), Fen-CTL vs. Fen-OTC (n = 3), Bog-CTL vs. Fen CTL, Bog-OTC vs. Fen-OTC (n = 3) and CTL vs. OTC (n = 6). Significant differences were tested using one-way ANOVA and are indicated by bold characters.

PLFAs	G+ bacteria	G– bacteria	Fungi	Actinobacteria	Protozoan
<i>Average value</i>					
CTL	19.5	7.3	2.7	6.7	5.7
OTC	29.8	11.5	6.1	9.5	5.9
Bog-CTL	14.1	5.7	1.4	3.9	2.3
Bog-OTC	12.3	4.2	6.0	7.2	4.3
Fen-CTL	25.0	8.8	4.1	9.5	9.0
Fen-OTC	47.3	18.8	6.1	11.7	7.5
<i>p-Value</i>					
CTL vs. OTC	0.38	0.37	0.28	0.41	0.91
Bog-CTL vs. Bog-OTC	0.89	0.74	0.43	0.48	0.49
Fen-CTL vs. Fen-OTC	0.14	0.14	0.59	0.65	0.18
Bog-CTL vs. Fen CTL	0.38	0.51	0.11	0.23	0.04
Bog-OTC vs. Fen-OTC	0.05	0.05	0.99	0.40	0.15
<i>Water content and WEOM features</i>					
	Water content	WEOC	Sugar content	SUVA ₂₈₀	
<i>Average value</i>					
CTL	93.8	2.9	1.2	0.066	
OTC	94.2	2.7	1.1	0.068	
Bog-CTL	93.3	2.9	1.2	0.073	
Bog-OTC	94.5	2.8	1.1	0.073	
Fen-CTL	94.3	2.9	1.1	0.059	
Fen-OTC	93.9	2.6	1.2	0.062	
<i>p-Value</i>					
CTL vs. OTC	0.27	0.38	0.81	0.84	
Bog-CTL vs. Bog-OTC	0.04	0.89	0.44	0.98	
Fen-CTL vs. Fen-OTC	0.40	0.32	0.92	0.62	
Bog-CTL vs. Fen CTL	0.07	0.84	0.77	0.23	
Bog-OTC vs. Fen-OTC	0.21	0.55	0.77	0.09	
<i>Extracellular enzymatic activities</i>					
	PO	BG	LAP	AP	
<i>Average value</i>					
CTL	0.0041	0.7	5.2	1.3	
OTC	0.0034	0.7	5.0	1.2	
Bog-CTL	0.0042	0.7	4.8	1.3	
Bog-OTC	0.0044	0.8	5.4	1.4	
Fen-CTL	0.0040	0.8	5.6	1.3	
Fen-OTC	0.0024	0.6	4.7	1.0	
<i>p-Value</i>					
CTL vs. OTC	0.45	0.84	0.72	0.60	
Bog-CTL vs. Bog-OTC	0.91	0.33	0.32	0.03	
Fen-CTL vs. Fen-OTC	0.13	0.14	0.10	0.13	
Bog-CTL vs. Fen CTL	0.75	0.34	0.29	0.84	
Bog-OTC vs. Fen-OTC	0.26	0.08	0.03	0.01	

4.2. Air warming can simultaneously lead to an increase in bacterial community in the fen and microbial activity in the bog

At the Bog site, air warming treatment led to higher AP enzymatic activity. This enzyme is produced by both soil microorganisms and plants and is involved in the mineralization of phosphate from phospholipids (Turner et al., 2002; Toor et al., 2003). AP changes underpin a higher breakdown of organically bound phosphate at the Bog site in the course of air warming. Particular attention must be paid to the impact of roots which are considered as key controlling factors of AP activity (Robroek et al., 2013). Indeed, air warming favoured vascular plant abundance rather than *Sphagnum* mosses (Jassey et al., 2013). Roots are lacking in *Sphagnum* species, and therefore it can be expected that

the root activity increase of vascular plants triggers AP activity. Jassey et al. (2013) also indicated that such a shift of vascular plants was associated with a decrease of *Sphagnum*-polyphenols, a strong microbial breakdown inhibitor, stimulating, in turn, bacterial and microbial enzymatic activities (Fenner and Freeman, 2011). Here, the lack of changes upon POA, BG and LAP provides no evidence for such a phenomenon.

Air warming also increased discrepancies between the Bog and the Fen sites when comparing first the control plots and then, the warmed plots of both sites. Thus, the increase in enzymatic activities at the warmed Bog site could be linked to higher temperature fluctuations in both the air and the soil, which may have triggered their kinetics (Davidson and Janssens, 2006). Additionally, PLFAs indicated that bacterial biomass increased in the warmed Fen site (Table 3), suggesting that air warming can alter the microbial structure and enzyme hydrolytic activities in opposite directions at the scale of the Bog and the Fen sites. Thus, air warming might induce the emergence of differential peat C dynamics in Bog and Fen sites. Following the soil C cycle scheme of Schimel and Weintraub (2003), one could hypothesize that C uptake by microbial cell biomass was favoured at the warmed Fen site, whereas it was hydrolytic enzyme production that was favoured at the warmed Bog site. However, it was also demonstrated in a snow removal experiment that differential timing of peat defrosting or snow melting can induce delays in the microbial community response (Robroek et al., 2013). Thus, the predominance of fungi upon bacterial biomass was used as an indicator of the winter state of the microbial community (Robroek et al., 2013). At the warmed Fen site, the relative shift of microbial structure to bacterial biomass could indicate that the microbial community was in a more advanced seasonal stage than at the Bog site. Moreover, an increase in enzymatic activities can also take place as a physiological adjustment to survive cold temperature (Beales, 2004). In any case, this advocates a more careful observation of the early spring period after snow melting.

Peat moisture was defined as the main controlling factor differentiating OM decomposition in the Bog and Fen sites (Delarue et al., 2011b). A change in moisture condition should therefore induce a change in peat C cycle. Here, we saw that water content did not differ in warmed plots nor in the control Bog and Fen plots. Such a change was strengthened by PLFAs from protozoa and indeed, it is known that testate amoebae are positively correlated to peat moisture and water-table depths (Woodland et al., 1998). The peat C cycle can be roughly divided into 4 components: soil organic C; dissolved organic C (WEOC in this study), microbial cell biomass and exoenzymes (Schimel and Weintraub, 2003). WEOC is an intermediate product between solid and gas phases in the course of decomposition (Schimel and Weintraub, 2003) and is considered as an indicator of the portion of dissolved OM which is the most active and mobile fraction within the OM (Akagi and Zsolnay, 2008; Zaccone et al., 2009). No significant change between control and warmed plots or between warmed Bog and Fen sites was recorded (Table 3). Moreover, WEOC mainly depended on sugar content which is known to be ubiquitous, occurring within both peat and microorganisms (results not shown). WEOC and sugar content were not discriminating enough to draw any conclusions about OM decomposition under the impact of OTCs in Bog and Fen sites. Additionally, since no change occurred upon SUVA₂₈₀, which should reflect the decomposition of recalcitrant aromatic moieties, our results suggest that air warming impacted neither recalcitrant nor labile OM pools in this study.

5. Conclusion

We have highlighted the limitation of the use of single environmental factors to assess the degree of OM decomposition. Instead, the use of multiple chemical, biological and microclimatic factors can provide more reliable information. For example, the single use of enzymatic activities would suggest that air warming indirectly favoured peat decomposition at the Bog sites, whereas our multidisciplinary approach

emphasises a change in microbial structure and activities as a consequence of complex interactions between groundwater level and air warming.

In addition, we have demonstrated that the enzymatic latch theory does not apply to cases where the water content does not decrease and consequently, where the oxygen level does not increase. This implies that a better definition of the environmental framework governing the enzymatic latch theory is needed, especially under future climatic conditions. Future investigations should aim at characterizing the seasonal pattern of these interactions, taking also into consideration soil microtopography, since this will greatly affect the impact of global warming on peat decomposition.

Acknowledgements

This work was funded as part of the PEATWARM initiative through an ANR (French National Agency for Research) grant (ANR-07-VUL-010). This paper is a contribution to the research conducted in the Labex VOLTAIRE (ANR-10-LABX-100-01). The authors are also indebted to the Regional Scientific Council of Natural Heritage of the Franche-Comté Region for access to Le Forbonnet site. They are grateful to CETRAHE laboratory for the analytical assistance. They would like to thank N. Pothier and L. Catherine for the logistical assistance and E. Rowley-Jolivet for the revision of the English version. The authors also thank the reviewers and Dr Mark Hanson for their constructive comments and helpful suggestions.

References

- Abbott, G.D., Swain, E.Y., Muhammad, A.B., Allton, K., Belyea, L.R., Laing, C.G., Cowie, G.L., 2013. Effect of water-table fluctuations on the degradation of *Sphagnum* phenols in surficial peats. *Geochim. Cosmochim. Acta* 106, 177–191.
- Akagi, J., Zsolnay, A., 2008. Effects of long-term de-vegetation on the quantity and quality of water extractable organic matter (WEOM): biogeochemical implications. *Chemosphere* 72, 1462–1466.
- Andersen, R., Grasset, L., Rochefort, L., Thormann, M.N., Francez, A.-J., 2010. Changes in microbial community structure and function following *Sphagnum* peatland restoration. *Soil Biol. Biochem.* 42, 291–301.
- Aronson, E.L., McNulty, S.G., 2009. Appropriate experimental ecosystem warming methods by ecosystem, objective, and practicality. *Agric. For. Meteorol.* 149, 1791–1799.
- Bardgett, R.D., Hobbs, P.J., Frostegård, Å., 1996. Changes in soil fungal:bacterial biomass ratios following reductions in the intensity of management of an upland grassland. *Biol. Fertil. Soils* 22, 261–264.
- Beales, N., 2004. Adaptation of microorganisms to cold temperatures, weak acid preservatives, low pH, and osmotic stress: a review. *Compr. Rev. Food Sci. Food Saf.* 3, 1–20.
- Bligh, E.G., Dyer, W.G., 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 37, 911–917.
- Bokhorst, S., Huiskes, A., Convey, P., Sinclair, B.J., Lebouvier, M., Van de Vijver, B., et al., 2011. Microclimate impacts of passive warming methods in Antarctica: implications for climate change studies. *Polar Biol.* 34, 1421–1435.
- Bokhorst, S., Huiskes, A., Aerts, R., Convey, P., Cooper, E.J., Dalen, L., et al., 2013. Variable temperature effects of Open Top Chambers at polar and alpine sites explained by irradiance and snow depth. *Glob. Chang. Biol.* 19, 64–74.
- Bragazza, L., Parisod, J., Buttler, A., Bardgett, R.D., 2013. Biogeochemical plant–soil microbe feedback in response to climate warming in peatlands. *Nat. Clim. Chang.* 3, 273–277.
- Breeuwer, A., Heijmans, M., Gleichman, M., Robroek, B.J.M., Berendse, F., 2009. Response of *Sphagnum* species mixtures to increased temperature and nitrogen availability. *Plant Ecol.* 204, 97–111.
- Carleton, T.J., Dunham, K.M.M., 2003. Distillation in a boreal mossy forest floor. *Can. J. For. Res.* 33, 663–671.
- Charman, D.J., Beilman, D.W., Blaauw, M., Booth, R.K., Brewer, S., Chambers, F.M., et al., 2013. Climate-related changes in peatland carbon accumulation during the last millennium. *Biogeosciences* 10, 929–944.
- Dabros, A., Fyles, J.W., 2010. Effects of open-top chambers and substrate type on biogeochemical processes at disturbed boreal forest sites in northwestern Quebec. *Plant Soil* 327, 465–479.
- Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440, 165–173.
- De Boeck, H.J., De Groot, T., Nijs, I., 2012. Leaf temperatures in glasshouses and open-top chambers. *New Phytol.* 194, 1155–1164.
- Delarue, F., Laggoun-Defarge, F., Buttler, A., Gogo, S., Jassey, V.E.J., Disnar, J.R., 2011a. Effects of short-term ecosystem experimental warming on water-extractable organic matter in an ombrotrophic *Sphagnum* peatland (Le Forbonnet, France). *Org. Geochem.* 42, 1016–1024.
- Delarue, F., Laggoun-Defarge, F., Disnar, J.R., Lottier, N., Gogo, S., 2011b. Organic matter sources and decay assessment in a *Sphagnum*-dominated peatland (Le Forbonnet, Jura Mountains, France): impact of moisture conditions. *Biogeochemistry* 106, 39–52.
- Dorrepaal, E., Toet, S., van Logtestijn, R.S.P., Swart, E., van de Weg, M.J., Callaghan, T.V., et al., 2009. Carbon respiration from subsurface peat accelerated by climate warming in the subarctic. *Nature* 460, 616–U79.
- Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A., Smith, F., 1956. Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 28, 350–356.
- Fenner, N., Freeman, C., 2011. Drought-induced carbon loss in peatlands. *Nat. Geosci.* 4, 895–900.
- Freeman, C., Liska, G., Ostle, N.J., Jones, S.E., Lock, M.A., 1995. The use of fluorogenic substrates for measuring enzyme-activity in peatlands. *Plant Soil* 175, 147–152.
- Freeman, C., Ostle, N., Kang, H., 2001. An enzymic 'latch' on a global carbon store – a shortage of oxygen locks up carbon in peatlands by restraining a single enzyme. *Nature* 409, 149–149.
- Freeman, C., Ostle, N.J., Fenner, N., Kang, H., 2004. A regulatory role for phenol oxidase during decomposition in peatlands. *Soil Biol. Biochem.* 36, 1663–1667.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., et al., 2006. Climate-carbon cycle feedback analysis: results from the (CMIP)-M-4 model intercomparison. *J. Clim.* 19, 3337–3353.
- Frostegård, Å., Bååth, E., 1996. The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biol. Fertil. Soils* 22, 59–65.
- Gorham, E., 1991. Northern peatlands – role in the carbon-cycle and probable responses to climatic warming. *Ecol. Appl.* 1, 182–195.
- Hansson, K., Kleja, D.B., Kalbitz, K., Larsson, H., 2010. Amounts of carbon mineralised and leached as DOC during decomposition of Norway spruce needles and fine roots. *Soil Biol. Biochem.* 42, 178–185.
- Hollister, R.D., Webber, P.J., Nelson, F.E., Tweedie, C.E., 2006. Soil thaw and temperature response to air warming varies by plant community: results from an open-top chamber experiment in northern Alaska. *Arct. Antarct. Alp. Res.* 38, 206–215.
- I.P.C.C. (Intergovernmental Panel on Climate Change), 2007. *Climate Change 2007: The Physical Science Basis*. Cambridge University Press.
- Jassey, V.E.J., Chiapusio, G., Gilbert, D., Buttler, A., Toussaint, M.L., Binet, P., 2011. Experimental climate effect on seasonal variability of polyphenol/phenoloxidase interplay along a narrow fen-bog ecological gradient in *Sphagnum fallax*. *Glob. Chang. Biol.* 17, 2945–2957.
- Jassey, V.E.J., Chiapusio, G., Binet, P., Buttler, A., Laggoun-Defarge, F., Delarue, F., et al., 2013. Above- and belowground linkages in *Sphagnum* peatland: climate warming affects plant–microbial interactions. *Glob. Chang. Biol.* 19, 811–823.
- Kalbitz, K., Schwesig, D., Schmerwitz, J., Kaiser, K., Haumaier, L., Glaser, B., et al., 2003. Changes in properties of soil-derived dissolved organic matter induced by biodegradation. *Soil Biol. Biochem.* 35, 1129–1142.
- Kroppenstedt, R.M., 1985. Fatty acid and menaquinone analysis of actinomycetes and related organisms. In: Goodfellow, M., Minnikin, D.E. (Eds.), *Chemical Methods in Bacterial Systematics*. Academic Press, London, pp. 173–199.
- Loisel, J., Yu, Z.C., 2013. Recent acceleration of carbon accumulation in a boreal peatland, south central Alaska. *J. Geophys. Res. Biogeosci.* 118, 41–53.
- Marion, G.M., Henry, G.H.R., Freckman, D.W., Johnstone, J., Jones, G., Jones, M.H., et al., 1997. Open-top designs for manipulating field temperature in high-latitude ecosystems. *Glob. Chang. Biol.* 3, 20–32.
- McLachley, G.P., Reddy, K.R., 1998. Regulation of organic matter decomposition and nutrient release in a wetland soil. *J. Environ. Qual.* 27, 1268–1274.
- McNeil, P., Waddington, J.M., 2003. Moisture controls on *Sphagnum* growth and CO₂ exchange on a cutover bog. *J. Appl. Ecol.* 40, 354–367.
- Otto, A., Otto, F.E.L., Boucher, O., Church, J., Hegerl, G., Forster, P.M., et al., 2013. Energy budget constraints on climate response. *Nat. Geosci.* 6, 415–416.
- Pind, A., Freeman, C., Lock, M.A., 1994. Enzymatic degradation of phenolic materials in peatlands – measurement of phenol oxidase activity. *Plant Soil* 159, 227–231.
- Price, J.S., Edwards, T.W.D., Yi, Y., Whittington, P.N., 2009. Physical and isotopic characterization of evaporation from *Sphagnum* moss. *J. Hydrol.* 369, 175–182.
- Ringelberg, D.B., Stair, J.O., Almeida, J., Norby, R.J., O'Neil, E.G., White, D.C., 1997. Consequences of rising atmospheric carbon dioxide levels for the belowground microbiota associated with white oak. *J. Environ. Qual.* 26, 495–503.
- Robroek, B.J.M., Heijboer, A., Jassey, V.E.J., Hefting, M.M.T., Rouwenhorst, G., Buttler, A., Bragazza, L., 2013. Snow cover manipulation effects on microbial community structure and soil chemistry in a mountain bog. *Plant Soil* 369, 151–164.
- Saiya-Cork, K.R., Sinsabaugh, R.L., Zak, D.R., 2002. The effects of long term nitrogen deposition on extracellular enzyme activity in an *Acer saccharum* forest soil. *Soil Biol. Biochem.* 34, 1309–1315.
- Schimel, J.P., Weintraub, M.N., 2003. The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biol. Biochem.* 35, 549–563.
- Sullivan, P.F., Arens, S.J.T., Chimner, R.A., Welker, J.M., 2008. Temperature and microtopography interact to control carbon cycling in a high arctic fen. *Ecosystems* 11, 61–76.
- Swain, E.Y., Abbott, G.D., 2013. The effect of redox conditions on sphagnum acid thermochemolysis product distributions in a northern peatland. *J. Anal. Appl. Pyrolysis* 103, 2–7.
- Toor, G.S., Condron, L.M., Di, H.J., Cameron, K.C., Cade-Menun, B.J., 2003. Characterization of organic phosphorus in leachate from a grassland soil. *Soil Biol. Biochem.* 35, 1317–1323.
- Turner, B.L., McKelvie, I.D., Haygarth, P.M., 2002. Characterisation of water-extractable soil organic phosphorus by phosphatase hydrolysis. *Soil Biol. Biochem.* 34, 27–35.
- van Breemen, N., 1995. How *Sphagnum* bogs down other plants. *Trends Ecol. Evol.* 10 (7), 270–275.

- Weedon, J.T., Kowalchuk, G.A., Aerts, R., van Hal, J., van Logtestijn, R., Tas, N., et al., 2012. Summer warming accelerates sub-arctic peatland nitrogen cycling without changing enzyme pools or microbial community structure. *Glob. Chang. Biol.* 18, 138–150.
- Weishaar, J.L., Aiken, G.R., Bergamaschi, B.A., Fram, M.S., Fujii, R., Mopper, K., 2003. Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Environ. Sci. Technol.* 37, 4702–4708.
- Wilkinson, S.C., 1988. Gram-negative bacteria. In: Ratledge, C., Wilkinson, S.C. (Eds.), *Microbial Lipids* vol. 1. Academic press, London, pp. 299–488.
- Woodland, W.A., Charman, D.J., Sims, P.C., 1998. Quantitative estimates of water tables and soil moisture in Holocene peatlands from testate amoebae. *The Holocene* 8, 261–273.
- Wretensjö, I., Svensson, L., Christie, W.W., 1990. Gas chromatographic–mass spectrometric identification of the fatty acids in borage oil using the picolinyl ester derivatives. *J. Chromatogr.* 521, 89–97.
- Yazaki, T., Urano, S., Yabe, K., 2006. Water balance and water movement in unsaturated zones of Sphagnum hummocks in Fuhrengawa Mire, Hokkaido, Japan. *J. Hydrol.* 319, 312–327.
- Zaccone, C., D'Orazio, V., Shotyky, W., Miano, T.M., 2009. Chemical and spectroscopic investigation of porewater and aqueous extracts of corresponding peat samples throughout a bog core (Jura Mountains, Switzerland). *J. Soils Sediments* 9, 443–456.
- Zelles, L., 1999. Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of microbial communities in soil: a review. *Biol. Fertil. Soils* 29, 111–129.